

New Skyrme Crystals

Paul Leask^{1,2}

Joint work with Derek Harland² & Martin Speight²

¹ p.n.leask@leeds.ac.uk

² School of Mathematics, University of Leeds

Geometric Models of Nuclear Matter, University of Kent, July 2022



UNIVERSITY OF LEEDS

Table of Contents

- 1 Motivation
- 2 Skyrme model
- 3 History of Skyrme crystals
- 4 Variational problem
- 5 Massless Skyrme crystals
- 6 Massive Skyrme crystals
- 7 Classical properties
- 8 Open problems



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems



Motivation



- The Skyrme model is a nonlinear field theory of pions (Skyrme, 1961)
- Nuclei are modelled as topological solitons (Skyrmions)
- Want to understand phases, and transitions of phases, of nuclear matter in the Skyrme model
- Ground state of nuclear matter has a crystalline structure in the classical approximation
- Many Skyrmions look like chunks of the infinite crystal (Feist *et al.*, 2013)
- Two candidates proposed:
 - Cubic lattice of half-Skyrmions
 - α -particle lattice
- Which is the lower energy solution (classically)?
- Are the two related?
- Are there other *new* solutions?



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

Skyrme model



- Topological solitons: smooth, spatially localized solutions of non-linear field theories, topologically stable against decay to vacuum.

- Skyrme field $\varphi : (\Sigma, g) \rightarrow (G, h)$, e.g. $\mathbb{R}^3/\Lambda \rightarrow \text{SU}(2)$

- Left-invariant Maurer-Cartan form $\theta = \varphi^{-1}d\varphi \in \Omega^1(G) \otimes \mathfrak{g}$
- Associated 2-form $\Omega \in \Omega^2(G) \otimes \mathfrak{g}$, $\Omega(X, Y) := [\theta(X), \theta(Y)]$
- $\text{Ad}(\text{SU}(2))$ invariant inner product, $h(X, Y) = \frac{1}{2} \text{Tr}(X^\dagger Y)$
- Skyrme energy functional

$$E[\varphi] = \int_{\Sigma} \left\{ c_2 g^{ij} h(L_i, L_j) + \frac{c_4}{2} g^{ia} g^{jb} h(\Omega_{ij}, \Omega_{ab}) + c_0 V(\varphi) \right\} \text{vol}_g$$

- $V : \text{SU}(2) \rightarrow \mathbb{R}$ is the pion mass potential,

$$V(\varphi) = m^2 \text{Tr}(\mathbb{1} - \varphi)$$

- Usual coupling constants $c_0 = c_2 = 1$ and $c_4 = 1/4$
- Derrick's scaling argument $E_4 = E_2 + 3E_0$
- Topological energy bound $E \geq 12\pi^2 \alpha |B|$ (Harland, 2014)
- Skyrmions are local minima of E
- Found numerically by discretising E and applying a gradient descent method



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

History of Skyrme crystals

History of Skyrme crystals



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals


Variational
problem

Massless
Skyrme
crystals


Massive
Skyrme
crystals

Classical
properties

Open
problems



(a) Half-Skyrmions lattice



(b) α -particle

- SC crystal of Skyrmions (Klebanov, 1985)
- BCC crystal of half Skyrmions (Goldhaber & Manton, 1987)
- SC crystal of half Skyrmions (Kugler & Shtrikman, 1988; Castillejo *et al.*, 1989)
- Building Skyrmions from the α -particle (Battye *et al.*, 2007)
- Massless (Silva Lobo, 2010) and massive (Adam *et al.*, 2022) phase transition between α -particle and SC crystal of half Skyrmions
- Constructing Skyrmions from crystal chunks (Feist *et al.*, 2013)
- Phase transitions between different crystals (Perapechka & Shnir, 2017)

History of Skyrme crystals




Figure: Skyrmions constructed from the α -particle

- (a) Constructed from two twisted α -particles (Battye *et al.*, 2007)
- (b) Constructed from three twisted α -particles (Battye *et al.*, 2007)
- (c) Constructed from six twisted α -particles (Feist *et al.*, 2013)
- (b) Constructed from eight α -particles (Battye *et al.*, 2007)

New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

Variational problem



Variational problem

- Warmup problem: Baby Skyrmion crystals (Leask, 2022)
- Skyrme crystals are maps

$$\varphi : \mathbb{R}^3/\Lambda \rightarrow \mathrm{SU}(2), \quad \Lambda = \{n_1 \mathbf{X}_1 + n_2 \mathbf{X}_2 + n_3 \mathbf{X}_3 : n_i \in \mathbb{Z}\}$$

- General idea (Speight, 2014): identify $(\mathbb{R}^3/\Lambda, \bar{g}) \longleftrightarrow (\mathbb{R}^3/\mathbb{Z}^3, g)$ via the diffeomorphism $F : \mathbb{T}^3 \rightarrow \mathbb{R}^3/\Lambda$ where $\mathbb{T}^3 = \mathbb{R}^3/\mathbb{Z}^3$ and $F(\mathbf{x}) = \mathcal{A}\mathbf{x}$, $\mathcal{A} = [\mathbf{X}_1 \ \mathbf{X}_2 \ \mathbf{X}_3]$
- Fix Skyrme field to be the map $\varphi : \mathbb{T}^3 \rightarrow \mathrm{SU}(2)$
- Metric on $\mathbb{T}^3$ is the pullback $g = F^*\bar{g}$, with $g_{ij} = \mathbf{X}_i \cdot \mathbf{X}_j$
- Vary metric g_s with $g_0 = F^*\bar{g} \iff$ vary lattice Λ_s with $\Lambda_0 = \Lambda$
- Energy minimized over all variations of $g \iff$ optimal period lattice Λ

New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems



Variational problem

- Let SPD_n be the space of symmetric positive-definite $n \times n$ -matrices.
- For fixed φ , can consider the Skyrme energy to be a map $E : \text{SPD}_3 \rightarrow \mathbb{R}$
- E is convex when restricted to geodesics $\Rightarrow$ uniqueness of the lattice
- Simple case: $\Lambda = L\mathbb{Z}^3$, energy scales as $E = LE_2 + \frac{1}{L}E_4 + L^3E_0$ and

$$\frac{dE}{dL} = 0 \quad \Rightarrow \quad L^2 = \frac{1}{2} \left(-\frac{E_2}{3E_0} + \sqrt{\left(\frac{E_2}{3E_0}\right)^2 + \frac{4E_4}{3E_0}} \right)$$

- In general, optimal period lattice Λ :
 - ◇ Massless pions: explicit solution
 - $\hookrightarrow$ Matrix square root
 - ◇ Massive pions: numerical solution
 - $\hookrightarrow$ Arrested Newton flow or nonlinear conjugate gradient descent

New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems



Matrix square root

- Massless case $V(\varphi) = 0$ reduces the problem to

$$\left(\frac{g\mathcal{K}}{\sqrt{\det g}} \right)^2 = \frac{c_2}{c_4} \mathcal{L}\mathcal{K},$$

where, in sigma model notation,

$$\mathcal{L}_{ij} = \int_{\Sigma} (\partial_i \pi \cdot \partial_j \pi) \text{vol}_g$$

and

$$\mathcal{K}^{ij} = \varepsilon^{iab} \varepsilon^{jcd} \int_{\Sigma} \{ (\partial_a \pi \cdot \partial_c \pi) (\partial_b \pi \cdot \partial_d \pi) - (\partial_a \pi \cdot \partial_d \pi) (\partial_b \pi \cdot \partial_c \pi) \} \text{vol}_g$$

- Matrix square root:

$$\frac{g\mathcal{K}}{\sqrt{\det g}} = PD^{1/2}P^{-1}, \quad D^{1/2} = \begin{pmatrix} \sqrt{\lambda_1} & 0 & 0 \\ 0 & \sqrt{\lambda_2} & 0 \\ 0 & 0 & \sqrt{\lambda_3} \end{pmatrix}$$

- Setting $\tilde{g} = \frac{g}{\sqrt{\det g}}$ yields $\tilde{g} = PD^{1/2}P^{-1}\mathcal{K}^{-1}$
- Finally, using the fact that $\det g = \frac{1}{(\det \tilde{g})^2}$, we arrive at $g = \frac{\tilde{g}}{\det \tilde{g}}$.



- Aim: solve the unconstrained optimisation problem

$$\min_{g \in \text{SPD}_3} E(g)$$

- Accelerated 2nd order gradient descent with flow arresting
- Solve Newton's equations of motion for a particle on SPD_3 with potential energy $E(g)$ using 4th order Runge–Kutta:

$$\partial_{ss} g_s|_{s=0} = - \left. \frac{\delta E}{\delta g_s} \right|_{s=0}, \quad g_0 = g$$

- Restart flow if $E(t + \delta t) > E(t)$ (arresting)
- Terminate flow when $\left. \frac{\delta E}{\delta g_s} \right|_{s=0} < 10^{-5}$ everywhere



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

Massless Skyrme crystals



1/2-lattice (FCC lattice)




Figure: Massless FCC lattice (Kugler & Shtrikman, 1988) and (Castillejo *et al.*, 1989)

- Obtained from Fourier series expansion initial configuration (Castillejo *et al.*, 1989)

$$\sigma = -c_1 c_2 c_3, \quad \pi^1 = s_1 \sqrt{1 - \frac{s_2^2}{2} - \frac{s_3^2}{2} + \frac{s_2^2 s_3^2}{3}}$$

and cyclic, where $c_i = \cos\left(\frac{2\pi x^i}{L}\right)$ and $s_i = \sin\left(\frac{2\pi x^i}{L}\right)$

- Symmetries (Kugler & Shtrikman, 1989):

$$A_1 : (x, y, z) \mapsto (-x, y, z)$$

$$(\sigma, \pi^1, \pi^2, \pi^3) \mapsto (\sigma, -\pi^1, \pi^2, \pi^3)$$

$$A_2 : (x, y, z) \mapsto (y, z, x)$$

$$(\sigma, \pi^1, \pi^2, \pi^3) \mapsto (\sigma, \pi^2, \pi^3, \pi^1)$$

$$C_3 : (x, y, z) \mapsto (x, z, -y)$$

$$(\sigma, \pi^1, \pi^2, \pi^3) \mapsto (\sigma, \pi^1, \pi^3, -\pi^2)$$

$$D_4 : (x, y, z) \mapsto (x + L/2, y, z)$$

$$(\sigma, \pi^1, \pi^2, \pi^3) \mapsto (-\sigma, -\pi^1, \pi^2, \pi^3)$$



New crystals from old

- Define a potential $V(Q) := E[Q\pi_{1/2}]$ for $Q \in \text{SO}(4)$
- Isospin symmetry group of the $1/2$ -lattice is

$$\Gamma = \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \right\}$$

- Then $V(QG) = V(Q) \forall G \in \Gamma$ and $V(SQ) = V(Q) \forall S \in \text{SO}(3)$
- So the potential is a map $V : \text{SO}(4)/\text{SO}(3) \cong S^3 \rightarrow \mathbb{R}$
- N/S-pole on S^3 corresponds to the $1/2$ -lattice
- Cubic symmetry group acts invariantly on equatorial S^2
- Intersection of S^2 with vertices, edge centres & face centres of the cube projected radially onto S^2 are critical points of V (principle of symmetric criticality)
- *Three* other Skyrme crystals should exist, one of which is the α -lattice

⇒ Two entirely new Skyrme crystals



- Let the map $SO(4)/SO(3) \cong S^3$ be given explicitly by

$$Q \mapsto Q^T \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \mathbf{v}_1, \quad Q^T = \begin{pmatrix} \uparrow & \uparrow & \uparrow & \uparrow \\ \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{v}_3 & \mathbf{v}_4 \\ \downarrow & \downarrow & \downarrow & \downarrow \end{pmatrix}$$

- Let $\{\mathbf{v}_1, \mathbf{u}_2, \mathbf{u}_3, \mathbf{u}_4\}$ be a set of linearly independent vectors in $\mathbb{R}^4$.
 - Use Gram-Schmidt to obtain an orthonormal basis $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ for $\mathbb{R}^4$
 - Can explicitly construct a Q such that $\pi = Q\pi_{1/2}$
- $\Rightarrow$ The three additional crystals can easily be obtained



4-lattice (α -lattice)




Figure: Massless α -particle lattice

- Obtained from the $B = 4$ rational map initial configuration (Houghton *et al.*, 1997)

$$R(z) = \frac{z^4 + 2\sqrt{3}iz^2 + 1}{z^4 - 2\sqrt{3}iz^2 + 1}$$

- Already shown the α -lattice and FCC lattice are related by an $SO(4)$ -isospin transformation (Leask (SIG X Talk), 2022):

$$Q\pi_4 = \pi_{1/2}, \quad Q = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 1/\sqrt{3} & -1/\sqrt{6} & 1/\sqrt{2} & 0 \\ 1/\sqrt{3} & -1/\sqrt{6} & -1/\sqrt{2} & 0 \\ 1/\sqrt{3} & 2/\sqrt{6} & 0 & 0 \end{pmatrix}$$



4-lattice (α -lattice)

- Related to the FCC lattice by $SO(4)$ -isospin transformation:

$$\pi_4 = Q\pi_{1/2}, \quad Q = \begin{pmatrix} 0 & -1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \\ 0 & 1/\sqrt{6} & -1/\sqrt{6} & 2/\sqrt{6} \\ 0 & -1/\sqrt{2} & -1/\sqrt{2} & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$$





Figure: Cube vertices projected radially onto S^2



(New) 1-lattice



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

- Related to the FCC lattice by $SO(4)$ -isospin transformation:

$$\pi_1 = Q\pi_{1/2}, \quad Q = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} & 0 \\ -1/\sqrt{6} & 2/\sqrt{6} & -1/\sqrt{6} & 0 \\ 1/\sqrt{2} & 0 & -1/\sqrt{2} & 0 \end{pmatrix}$$





Figure: Cube face centres projected radially onto S^2



(New) 2-lattice



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

- Related to the FCC lattice by $SO(4)$ -isospin transformation:

$$\pi_2 = Q\pi_{1/2}, \quad Q = \begin{pmatrix} 0 & 0 & 1/\sqrt{2} & 1/\sqrt{2} \\ 1 & 0 & 0 & 0 \\ 0 & 2/\sqrt{6} & -1/\sqrt{6} & 1/\sqrt{6} \\ 0 & 1/\sqrt{3} & 1/\sqrt{3} & -1/\sqrt{3} \end{pmatrix}$$





Figure: Cube edge centres projected radially onto S^2



Massless Skyrme crystals



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems




Figure: Skyrme crystals for $m = 0$. Top row is baryon density. Bottom row is sigma plots, where $\sigma = 0.9$ and $\sigma = -0.9$.



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

Massive Skyrme crystals

Massive Skyrme crystals



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals


Variational
problem

Massless
Skyrme
crystals


Massive
Skyrme
crystals

Classical
properties


Open
problems




(a) 1/2-lattice



(b) 4-lattice



(c) 1-lattice



(d) 2-lattice







Figure: Skyrme crystals for $m = 1$. Top row is baryon density. Bottom row is sigma plots, where $\sigma = 0.9$ and $\sigma = -0.9$.



Isospin moment of inertia tensor

- Collective coordinate approach to isospin d.o.f. (Adkins *et al.*, 1983)

$$\varphi(x) \mapsto \hat{\varphi}(x, t) = A(t)\varphi(x)A^\dagger(t).$$

Isorotations are symmetries of E so these configurations are all energy-degenerate.

- Isorotational angular velocity is $\omega_j = -i \text{Tr}(\tau^j A^\dagger \dot{A})$
- Maurer-Cartan form transforms as

$$\hat{L}_\mu = \hat{\varphi}^\dagger \partial_\mu \hat{\varphi} = \begin{cases} A\omega_i T_i A^\dagger, & \mu = 0 \\ AL_i A^\dagger, & \mu = i = 1, 2, 3. \end{cases}$$

- $T_i = \frac{i}{2} \varphi^\dagger [\tau^i, \varphi]$ is also an $\mathfrak{su}(2)$ current
- Effective Lagrangian on restricted space of configurations is $L_{\text{eff}} = L_{\text{rot}} - M_B$, where

$$L_{\text{rot}} = \frac{1}{2} \omega_i U_{ij} \omega_j$$

and the isospin moment of inertia is

$$U_{ij} = - \int_{\Sigma} \text{Tr} \left(c_2 T_i T_j + c_4 g^{ab} [L_a, T_i] [L_b, T_j] \right) \text{vol}_g$$



New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

Classical properties

Massless Skyrme crystals





4-lattice	1/2-lattice
	
$E^4 = 1.0378$	$E^{1/2} = 1.0378$
$U^4 = \begin{pmatrix} 238.6 & 0 & 0 \\ 0 & 238.6 & 0 \\ 0 & 0 & 297.9 \end{pmatrix}$	$U^{1/2} = \begin{pmatrix} 297.9 & 0 & 0 \\ 0 & 297.9 & 0 \\ 0 & 0 & 297.9 \end{pmatrix}$

Table: Comparison of the massless ($m = 0$) 4-lattice and 1/2-lattice

New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

New massless Skyrme crystals





1-lattice	2-lattice
	
$E^1 = 1.0378$	$E^2 = 1.0378$
$U^1 = \begin{pmatrix} 238.6 & 0 & 0 \\ 0 & 238.6 & 0 \\ 0 & 0 & 297.9 \end{pmatrix}$	$U^2 = \begin{pmatrix} 238.6 & 0 & 0 \\ 0 & 238.6 & 0 \\ 0 & 0 & 297.9 \end{pmatrix}$

Table: Comparison of the massless ($m = 0$) 1-lattice and 2-lattice

New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

Massive Skyrme crystals





4-lattice	1/2-lattice
	
$E^4 = 1.0631$	$E^{1/2} = 1.0673$
$U^4 = \begin{pmatrix} 135.5 & 0 & 0 \\ 0 & 135.5 & 0 \\ 0 & 0 & 167.3 \end{pmatrix}$	$U^{1/2} = \begin{pmatrix} 165.2 & 0 & 0 \\ 0 & 165.2 & 0 \\ 0 & 0 & 165.2 \end{pmatrix}$

Table: Comparison of the massive ($m = 1$) 4-lattice and 1/2-lattice

New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems

New massive Skyrme crystals





1-lattice	2-lattice
	
$E^1 = 1.0629$	$E^2 = 1.0631$
$U^1 = \begin{pmatrix} 166.8 & 0 & 0 \\ 0 & 135.8 & 0 \\ 0 & 0 & 135.8 \end{pmatrix}$	$U^2 = \begin{pmatrix} 167.2 & 0 & 0 \\ 0 & 135.7 & 0 \\ 0 & 0 & 135.6 \end{pmatrix}$

Table: Comparison of the massive ($m = 1$) 1-lattice and 2-lattice

New
Skyrme
Crystals

Paul Leask

Motivation

Skyrme
model

History of
Skyrme
crystals

Variational
problem

Massless
Skyrme
crystals

Massive
Skyrme
crystals

Classical
properties

Open
problems



Comparison of the four crystals

m	E^4	$E^{1/2}$	E^1	E^2
0	1.0378	1.0378	1.0378	1.0378
1	1.0631	1.0673	1.0629	1.0631
3	1.0710	1.0797	1.0703	1.0708
5	1.0715	1.0816	1.0707	1.0713
10	1.0710	1.0824	1.0703	1.0707

Table: Comparison of the four crystals for various m

- Massless crystals:
 - Energy degenerate
 - Isospin tensors share common eigenvalue(s)
 - All related to one another by $SO(4)$ -isospin transformations
- Massive crystals:
 - Energy is not invariant under $SO(4)$ action
 - ⇒ New lower energy crystal – the 1-lattice



Open problems



- What happens when we consider quantum corrections from the isospin d.o.f. for these two new crystals? (Adam *et al.*, 2022) investigated this for the 1/2-lattice and the 4-lattice.
- How do the energies of the new 1-lattice and 2-lattice compare to the 4-lattice at high and low density (e.g. produce the $E(V)$ curve)?
 - New lattices are not cubic, do they scale with the pion mass?
- How can we construct Skyrmions from these new lattices?